

Logarithmic corrections to black hole and black ring entropy in tunneling approach

Yi-Xin Chen

Zhejiang Institute of Modern Physics, Zhejiang University, Hangzhou, 310027, China
E-mail: yxchen@zimp.zju.edu.cn

Jian-Long Li

Zhejiang Institute of Modern Physics, Zhejiang University, Hangzhou, 310027, China
E-mail: marryrene@gmail.com

Kai-Nan Shao

Zhejiang Institute of Modern Physics, Zhejiang University, Hangzhou, 310027, China
E-mail: shaokn@gmail.com

ABSTRACT: The tunneling approach beyond semiclassical approximation has been used to calculate the corrected Hawking temperature and entropy for various black holes and FRW universe model. We examine their derivations, and prove that the quantity H in the corrected temperature is the explicit function of the only free parameter $\mathcal{A}$ (which is an auxiliary parameter defined by $\mathcal{A} = \hbar S_{BH}$). Our analysis improves previous calculations, and indicates that the leading order logarithmic correction to entropy is a natural result of the corrected temperature and the first law of thermodynamics. Additionally, we apply the tunneling approach beyond semiclassical approximation to neutral black rings. Based on the analysis, we show that the entropy of neutral black rings also has a logarithmic leading order correction.

KEYWORDS: GR black holes, gravity, quantum field theory on curved space .

Contents

1. Introduction	1
2. Exact differential and logarithmic correction to black hole entropy	3
3. Corrected entropy of neutral black rings	7
4. Summary	9

1. Introduction

Since the discovery of Hawking radiation and black hole thermodynamics[1, 2, 3], it is generally believed that black holes are objects which have temperature and entropy. By adopting the Hawking temperature $T_H = \frac{\kappa}{2\pi}$ and the Bekenstein-Hawking entropy $S_{BH} = \frac{A_H}{4\hbar}$, the first law of black hole thermodynamics is obtained[4]. The understanding of black hole entropy is an important subject in quantum gravity. When quantum effects are considered, the area law of black hole entropy should undergo corrections. These corrections have been extensively studied using many approaches, such as field theory methods[5, 6, 4], quantum geometry techniques[7, 8], general statistical mechanical arguments[9, 10, 11], Cardy formula[12, 13]. In all these derivations, the leading order correction to black hole entropy takes a logarithmic form

$$S = S_{BH} + \alpha \ln S_{BH} + \cdots . \quad (1.1)$$

This result also coincides with that obtained by counting the number of microstates in string theory[14, 15] and loop quantum gravity[16, 17].

The simplest method of calculating black hole entropy is directly integrating from the the first law of thermodynamics $dM = TdS$, using the expression of Hawking temperature. Several derivations of Hawking radiation were presented[18, 19, 20], one of which is the tunneling approach[21, 22]. In this approach, the emission rate of a particle tunneling from the black hole is associated with the imaginary part of the action, which in turn is related to the Boltzmann factor for the emission at the Hawking temperature. There are two variants of the tunneling approach, namely, Parikh-Wilczek radial null geodesic method[22, 23], and the Hamilton-Jacobi method[21, 24, 25]. Recently, based on the Hamilton-Jacobi method, Banerjee and Majhi[26, 27] developed a general formalism of tunneling beyond semiclassical approximation. According to their calculations, the Hawking temperature with quantum corrections takes the form

$$T_{cr} = T_H \left(1 + \sum \gamma_i \hbar^i \right) = T_H \left(1 + \sum \beta_i \frac{\hbar^i}{H^i} \right)^{-1} , \quad (1.2)$$

where H is a quantity with the dimension of $\hbar$, and should be expressed in terms of black hole parameters[28]. It is reasonable to assume that, the first law of thermodynamics still holds in the context of quantum corrections, and the entropy S_{cr} is a state function. Then, the entropy can be determined by integrating the relation $dE = T_{cr}dS_{cr}$, which has a form as (1.1). This formulism has been applied on various black holes[29, 28, 30, 31, 32] and FRW universe model[33]. However, after examining their papers carefully, we find that these derivations rely on one procedure, i.e., they expressed H as a polynomial of selected black hole parameters. In fact, all what we know is that H can be expressed in terms of black hole parameters. There is no reason why H “should” be a polynomial, and why H took those different particular forms in various models[29, 28, 30, 31, 32]. In order to fill this gap in the calculations of the quantum corrected entropy using the tunneling approach beyond semiclassical approximation, we propose to take H as a general function of a complete set of black hole parameters, i.e. $H = H(\text{black hole parameters})$. For example, in Kerr-Newman black holes, the complete parameters are M , J , and Q , while the entropy S_{BH} can be expressed in terms of them $S_{BH} = S_{BH}(M, Q, J)$. Next we define an auxiliary $\mathcal{A}$ by $\mathcal{A} := \mathcal{A}(M, J, Q) = \hbar S_{BH}$. This is a quantity with area unit, and can still be defined via the above formula even when the entropy S_{BH} does not obey the Bekenstein-Hawking area law¹. We may then regard the parameters (J, Q, S_{BH}) , or equivalently $(J, Q, \mathcal{A})$ as a complete set of parameters for Kerr-Newman black hole. So H can be expressed as $H = H(M, Q, J) = H(J, Q, \mathcal{A})$. According to the spirit in [28], by demanding dS_{cr} to be an exact differential, we can prove that $\frac{\partial H(J, Q, \mathcal{A})}{\partial J} = 0 = \frac{\partial H(J, Q, \mathcal{A})}{\partial Q}$, and so $H = H(\mathcal{A})$. Then performing the dimension analysis as in [29, 28, 30, 31, 32], we can show that the leading order correction to black hole entropy is logarithmic. Comparing to previous calculations, the treatment here is more natural and general. In particular, this treatment emphasizes the use of exact differential in eliminating the extra dependence in H . Another advantage is that the analysis here can be easily extended to the cases of AdS Schwarzschild black holes, Gauss-Bonnet and Lovelock black holes, the FRW universe model and neutral black rings. To this context, we can say that our analysis indicates that the leading order logarithmic correction to black hole entropy may be a general result naturally obtained from the corrected temperature (1.2) and the first law of thermodynamics.

The 5-dimensional black ring is found as a vacuum solution of general relativity, with the event horizon topology $S^1 \times S^2$ [34]. It is very interesting to study the corrected temperature and entropy of black rings due to quantum effects. The tunneling formulism can be used to examine the Hawking radiation of black rings[35, 36]. However, the correction due to quantum effects is not considered. On the other hand, according to[37], the near horizon dynamics of a scalar field φ on black ring metric background can be reduced to a $(1+1)$ dimensional one. Therefore, the tunneling formalism beyond semiclassical approximation can also be applied to black rings. In this Letter, we perform the analysis on neutral black ring, and show that its corrected temperature and entropy should also take

¹One example of violating the area law is the Lovelock gravity, for consistency, in this situation we still use S_{BH} to denote the standard black hole entropy without quantum correction. Another point is that one should not confuse $\mathcal{A}$ with the horizon area of black holes. We shall denote the horizon area by A_H , if appears in this paper.

the form of (1.2) and (1.1). As far as we know, it is the first time to propose that the entropy of neutral black ring should also undergoes a logarithmic leading order correction due to quantum effects. In obtaining the corrected entropy from the temperature (1.2), the previous treatment in [29, 28, 30, 31, 32] seems does not work with black rings. The parameters for neutral black ring are R , ν and λ (with $\lambda = 2\nu/(1 + \nu^2)$), while only R is dimensional. We are not able to determine (or ‘guess’) a particular polynomial for H . However, we may regard (R, ν) , (M, J) or equivalently $(J, \mathcal{A})$ as a complete set of parameters for neutral black rings. Then our treatment for Kerr-Newman can easily be extended on the case of black rings. Although the black hole and black ring entropy can only be fully determined by counting the microstate in well defined quantum gravity theories, the study of corrected entropy due to quantum effects would be helpful for the understanding of these black objects.

The paper is organized as follows. In Section 2, we prove that in the quantity H in the tunneling formalism beyond semiclassical approximation is an explicit function of the only parameter $\mathcal{A}$. We illustrate our proof in the context of Kerr-Newman black holes, and show that this analysis can easily be extended to the cases of AdS Schwarzschild black holes, Gauss-Bonnet and Lovelock black holes, and FRW universe model. In Section 3, we apply the tunneling formulism on neutral black ring, and obtain the corrected temperature and entropy. We use the units $G = c = k_B = \frac{1}{4\pi\epsilon_0} = 1$, while keeping the Planck constant $\hbar$ explicitly. In fact, the Planck constant is an indication of quantum effects, which can be used to identify the leading order correction.

2. Exact differential and logarithmic correction to black hole entropy

In this section, we take the Kerr-Newman black hole as an example to demonstrate that the leading order correction to the entropy is logarithmic, and then extend our treatment to other models. Before calculating the entropy correction from the corrected Hawking temperature (1.2) obtained in [26, 27], we give a brief review of the first law of thermodynamics on Kerr-Newman black holes. The standard first law holds on the event horizon

$$dS_{BH} = \frac{dM}{T_H} + \left(-\frac{\Omega_H}{T_H}\right) dJ + \left(-\frac{\Phi_H}{T_H}\right) dQ, \quad (2.1)$$

where Ω_H and Φ_H are the angular velocity and the electrical potential on the horizon. In Eq.(2.1), the black hole can be treated as an equilibrium state parametrized by M , J and Q , and the entropy S_{BH} is a state function. Integrate Eq.(2.1) by M , J and Q , S_{BH} can be expressed as a function of these quantities. For example, the Bekenstein-Hawking entropy of Kerr-Newman black hole takes the form

$$S_{BH} = \frac{\pi}{\hbar} \left(2M \left[M + \sqrt{M^2 - \frac{J^2}{M^2} - Q^2} \right] - Q^2 \right). \quad (2.2)$$

The thermodynamical quantities T_H , Ω_H and Φ_H can be defined as partial derivatives of

S_{BH}

$$\left. \frac{\partial S_{BH}}{\partial M} \right|_{J,Q} = \frac{1}{T_H} , \quad (2.3)$$

$$\left. \frac{\partial S_{BH}}{\partial J} \right|_{M,Q} = -\frac{\Omega_H}{T_H} , \quad (2.4)$$

$$\left. \frac{\partial S_{BH}}{\partial Q} \right|_{M,J} = -\frac{\Phi_H}{T_H} . \quad (2.5)$$

Besides, S_{BH} is a state function and dS_{BH} is an exact differential. This leads to the following Maxwell's relations

$$\left. \frac{\partial}{\partial J} \left(\frac{1}{T_H} \right) \right|_{M,Q} = \left. \frac{\partial}{\partial M} \left(-\frac{\Omega_H}{T_H} \right) \right|_{J,Q} , \quad (2.6)$$

$$\left. \frac{\partial}{\partial Q} \left(-\frac{\Omega_H}{T_H} \right) \right|_{M,J} = \left. \frac{\partial}{\partial J} \left(-\frac{\Phi_H}{T_H} \right) \right|_{M,Q} , \quad (2.7)$$

$$\left. \frac{\partial}{\partial M} \left(-\frac{\Phi_H}{T_H} \right) \right|_{J,Q} = \left. \frac{\partial}{\partial Q} \left(\frac{1}{T_H} \right) \right|_{J,M} . \quad (2.8)$$

When $\hbar \rightarrow 0$, the corrected temperature $T_{cr} \rightarrow T_H$, so it is reasonable to assume that the first law of thermodynamics still holds in the context of quantum correction[28], i.e.

$$dS_{cr} = \frac{dM}{T_{cr}} + \left(-\frac{\Omega_H}{T_{cr}} \right) dJ + \left(-\frac{\Phi_H}{T_{cr}} \right) dQ .$$

The corrected Hawking temperature calculated from the tunneling formalism beyond the semiclassical approximation is

$$T_{cr} = T_H \left(1 + \sum \gamma_i \hbar^i \right) = T_H \left(1 + \sum \beta_i \frac{\hbar^i}{H_{KN}^i} \right)^{-1} , \quad (2.9)$$

where H_{KN} is a quantity with dimension $\hbar$. Similar Maxwell's relations hold by replacing T_{BH} with T_{cr} . By substituting the expression of corrected temperature (2.9), we obtain the corresponding constrain equations for H_{KN}

$$\left. \frac{\partial H_{KN}}{\partial J} \right|_{M,Q} = -\Omega_H \left. \frac{\partial H_{KN}}{\partial M} \right|_{J,Q} , \quad (2.10)$$

$$\left. \frac{\partial H_{KN}}{\partial Q} \right|_{M,J} = \left(\frac{\Phi_H}{\Omega_H} \right) \left. \frac{\partial H_{KN}}{\partial J} \right|_{M,Q} , \quad (2.11)$$

$$\left. \frac{\partial H_{KN}}{\partial M} \right|_{J,Q} = -\frac{1}{\Phi_H} \left. \frac{\partial H_{KN}}{\partial Q} \right|_{J,M} . \quad (2.12)$$

In Kerr-Newman black hole, the three conserved quantities M , J , and Q have totally determined the black hole thermodynamics, so the quantity H_{KN} should be expressed in terms of these black hole parameters. The next step in [28] is to express H_{KN} as a particular polynomial of M, J, Q , followed by a tedious analysis using the first law of thermodynamics

to determine the expression of H_{KN} . This treatment is not natural and general, and can be improved. Here we take the general form $H_{KN} = H_{KN}(M, J, Q)$. Next we introduce an auxiliary parameter $\mathcal{A}$ defined by $\mathcal{A} = \hbar S_{BH}$. Then we can take $\mathcal{A}$ and two of the three conserved quantities M, J, Q as free parameters, and construct the third conserved quantity as a function of them. For example, taking $M = M(\mathcal{A}, J, Q)$ and substituting it into H_{KN} , we get

$$H_{KN} = H_{KN}(M, J, Q) = H'_{KN}(\mathcal{A}(M, J, Q), J, Q) .$$

Here we denote $H'_{KN}(\mathcal{A}, J, Q)$ again as $H_{KN}(\mathcal{A}, J, Q)$ for convenience. Substitute $H_{KN}(\mathcal{A}(M, J, Q), J, Q)$ into the constrain equation (2.10), and use the chain rule in standard calculus textbooks,

$$\begin{aligned} LHS. = & \frac{\partial H_{KN}(\mathcal{A}, J, Q)}{\partial \mathcal{A}} \bigg|_{J, Q} \frac{\partial \mathcal{A}}{\partial J} \bigg|_{M, Q} \\ & + \frac{\partial H_{KN}(\mathcal{A}, J, Q)}{\partial Q} \bigg|_{\mathcal{A}, Q} \frac{\partial Q}{\partial J} \bigg|_{M, Q} + \frac{\partial H_{KN}(\mathcal{A}, J, Q)}{\partial J} \bigg|_{\mathcal{A}, Q} , \end{aligned} \quad (2.13)$$

$$\begin{aligned} RHS. = & -\Omega_H \left(\frac{\partial H_{KN}(\mathcal{A}, J, Q)}{\partial \mathcal{A}} \bigg|_{J, Q} \frac{\partial \mathcal{A}}{\partial M} \bigg|_{J, Q} \right. \\ & \left. + \frac{\partial H_{KN}(\mathcal{A}, J, Q)}{\partial J} \bigg|_{\mathcal{A}, Q} \frac{\partial J}{\partial M} \bigg|_{J, Q} + \frac{\partial H_{KN}(\mathcal{A}, J, Q)}{\partial Q} \bigg|_{\mathcal{A}, J} \frac{\partial Q}{\partial M} \bigg|_{J, Q} \right) . \end{aligned} \quad (2.14)$$

The second term of Eq.(2.13), the second and the third term of Eq.(2.14) vanish. In addition, the first term of Eq.(2.13) cancels with the first term of Eq.(2.14) due to Eq.(2.4). Thus we obtain one constraint of H_{KN}

$$\frac{\partial H_{KN}(\mathcal{A}, J, Q)}{\partial J} \bigg|_{\mathcal{A}, Q} = 0 . \quad (2.15)$$

Similarly, the other constraint of H_{KN} can be obtained from the constraint equation (2.12)

$$\frac{\partial H_{KN}(\mathcal{A}, J, Q)}{\partial Q} \bigg|_{\mathcal{A}, J} = 0 . \quad (2.16)$$

As a result, once we take $\mathcal{A}$ as a free parameter in describing H_{KN} , it is the only explicit free parameter, i.e.

$$H_{KN} = H_{KN}(\mathcal{A}) . \quad (2.17)$$

If we take $J = J(\mathcal{A}, M, Q)$ or $Q = Q(\mathcal{A}, M, J)$, Eq.(2.17) can also be obtained from the constraint equations (2.10-2.12). This indicates that the result (2.17) is a naturally consequence of the first law of thermodynamics. Now we have completed the proof that H_{KN} is the function of the only explicit parameter $\mathcal{A}$.

In D-dimensional spacetime, the units $G = c = k_B = \frac{1}{4\pi\epsilon_0} = 1$, and $[H_{KN}] = [\hbar] = [\mathcal{A}] = L^{D-2}$, so in general H_{KN} can be expressed as²

$$H_{KN} = \eta \mathcal{A} , \quad (2.18)$$

²One can expect that the dependence of $\hbar$ in T_{cr} has been expressed in (2.9), so that $\hbar$ does not appear in the expression of H_{KN} .

where η is a dimensionless parameter. Using the first law of thermodynamics

$$\begin{aligned} dS_{cr} &= \frac{1}{T_{cr}} (dM - \Omega_H dJ - \Phi_H dQ) \\ &= \frac{1}{T_{cr}} (T_H dS_{BH}) , \end{aligned}$$

which can be written as

$$\begin{aligned} dS_{cr} &= \left(1 + \sum \beta_i \frac{\hbar^i}{H_{KN}^i} \right) dS_{BH} \\ &= \left(1 + \beta_1 \frac{\hbar}{\eta \mathcal{A}} + O(\hbar^2) \right) dS_{BH} \\ &= \left(1 + \beta_1 \frac{1}{\eta S_{BH}} + O(\hbar^2) \right) dS_{BH} \\ &= \left(1 + \frac{\alpha}{S_{BH}} + \dots \right) dS_{BH} \end{aligned} \tag{2.19}$$

where α is another dimensionless coefficient, and all other higher order terms are included in $(\dots)$. Integrate Eq.(2.19) up to the first (leading) order, which yields

$$S_{cr} = S_{BH} + \alpha \ln S_{BH} + \dots + const . \tag{2.20}$$

This is the corrected entropy of Kerr-Newman black hole obtained from the temperature (2.9) and the leading order correction is logarithmic.

Our analysis can be easily extended to the cases of various black holes and FRW universe model[28, 29, 30, 31, 32, 33], without assigning H as specific polynomials for each case :

- For Schwarzschild black hole , the only conserved quantity is mass M , and $S_{BH} = \frac{A_H}{4\hbar} = \frac{4\pi M^2}{\hbar} \equiv \frac{A}{\hbar}$ so $H = H(M) = H(\mathcal{A})$. The cases of Kerr black hole and Reissner-Nordstrom black hole is similar to the Kerr-Newman black hole.
- For FRW universe model, the parameter is $\tilde{r}_A$, the radius of apparent horizon, and $S_{BH} = \frac{\pi \tilde{r}_A^2}{\hbar} \equiv \frac{A}{\hbar}$. So $H(\tilde{r}_A) = H(\mathcal{A})$.
- For AdS Schwarzschild black hole, the situation is a bit more complicated. The mass and entropy of AdS Schwarzschild black hole is

$$M = \frac{r_+(1 + r_+^2/l^2)}{2} , \quad S = \frac{\pi r_+^2}{\hbar} ,$$

where $\Lambda = -3l^2$, l is the AdS radius, and $[l]^2 = L^2 = [\hbar]$. Since the cosmological constant comes into the black hole solution, it modifies the form of first law of black hole thermodynamics. Thus it is reasonable to consider an ensemble where the cosmological constant can fluctuate[38],

$$dS = \frac{1}{T} (dM - \Omega dJ - \Phi dQ - \Theta d\Lambda). \tag{2.21}$$

Then we can prove that $H = H(\mathcal{A})$. This result not only preserves the consistency of the black hole thermodynamics in AdS case, but also ensures that when $\Lambda \rightarrow 0$, the quantum corrections of black hole temperature and entropy degenerate to the Schwarzschild case.

- For D-dimensional Gauss-Bonnet black hole ,

$$S_{GB} = \frac{A_{D-2} r_h^{D-2}}{4\hbar} \left(1 + \frac{D-2}{D-4} \frac{\bar{\lambda}}{r_h^2} \right) ,$$

where $\bar{\lambda} = \lambda(D-3)(D-4)$, $[\bar{\lambda}] = L^2$, and λ is the coupling constant of the Gauss-Bonnet term (see [31]). It undergoes similar situation as the AdS case, and we can also get $H = H(\mathcal{A})$.

There is one comment on the corrected entropy of AdS Schwarzschild black hole. In the treatment of [26], they set $H = M^2$ for the AdS Schwarzschild black hole (See Eq.(4.9) and (4.10) therein). Following their treatment, we can write the terms of quantum corrections to the entropy explicitly as

$$S_{cr} - S_{BH} = \ln S_{BH} + \frac{\pi l^2}{\pi l^2 + \hbar S_{BH}} - \ln \left(\frac{\pi l^2}{\hbar} + S_{BH} \right) + \dots \quad . \quad (2.22)$$

When the cosmological constant vanishes, i.e., as $\Lambda = -3l^{-2} \rightarrow 0$, the dynamics of AdS Schwarzschild black hole should reduce to the corresponding asymptotically flat Schwarzschild black hole. However, as $\Lambda \rightarrow 0$, the above correction (2.22) is not logarithmic, which does not coincide with the the corrected entropy of Schwarzschild black hole. So the treatment in [26] cannot be correct. After examining their derivation, we think that the choice of $H = M^2$ for AdS Schwarzschild black hole is groundless and dogmatic. Their result in Gauss-Bonnet black hole suffers similar problems. In our paper, we can overcome this arbitrariness, and obtain the corrected leading order logarithmic correction to black hole entropy.

Now we have examined the derivations of corrected entropy in the cases of these various black holes [29, 28, 30, 31, 32], FRW universe model[33], and added the proof of the expression $H = H(\mathcal{A})$. Our treatment strengthens the validity of the method of obtaining the corrected entropy from the tunneling formalism beyond semiclassical approximation. The analysis indicate that the leading order entropy correction is a natural result obtained from the corrected temperature (2.9) and the first law of thermodynamics. In the next section we shall extend the above analysis to neutral black rings, and shows that its leading order entropy correction due to quantum effects is also logarithmic.

3. Corrected entropy of neutral black rings

The 5-dimensional black ring is a vacuum solution of general relativity, with the event horizon topology $S^1 \times S^2$. The tunneling approach has also been used to analyze the black ring radiation[35, 36]. However, the influence of quantum effects are not included. In

this section we apply the tunneling method beyond semiclassical approximation on neutral black rings. The metric of neutral black ring is

$$ds^2 = -\frac{F(y)}{F(x)}(dt - CR\frac{1+y}{F(y)}d\psi)^2 + \frac{R^2}{(x-y)^2}F(x) \\ \times \left[-\frac{G(y)}{F(y)}d\psi^2 - \frac{dy^2}{G(y)} + \frac{dx^2}{G(x)} + \frac{G(x)}{F(x)}d\phi^2 \right]$$

where $F(\xi) = 1 + \lambda\xi$, $G(\xi) = (1 - \xi^2)(1 + \nu\xi)$, $C(\nu, \lambda) = \sqrt{\lambda(\lambda - \nu)\frac{1+\lambda}{1-\lambda}}$. λ, ν are dimensionless parameters taking values in the range $0 < \nu \leq \lambda < 1$. x is one of the angular coordinates in S^2 , with $-1 \leq x \leq 1$. y is only “ringlike-radical” coordinate, with $-\infty \leq y \leq -1$. To remove the conical singularity at $x = 1$, λ and ν must be related to each other via

$$\lambda = \frac{2\nu}{1 + \nu^2} . \quad (3.1)$$

The horizon is at $y = y_h = -1/\nu$ with the topology $S^1 \times S^2$.

The thermodynamics quantities of neutral black ring are

$$M = \frac{3\pi R^2}{4} \frac{\lambda}{1 - \nu} , \quad (3.2)$$

$$J = \frac{\pi R^3}{2} \frac{\sqrt{\lambda(\lambda - \nu)(1 + \lambda)}}{(1 - \nu)^2} , \quad (3.3)$$

$$S_{\text{BR}} = \frac{\mathcal{A}_h}{4\hbar} = \frac{2\pi^2 R^3}{\hbar} \frac{\nu^{3/2} \sqrt{\lambda(1 - \lambda^2)}}{(1 - \nu)^2(1 + \nu)} , \quad (3.4)$$

$$T_{\text{BR}} = \frac{\hbar}{4\pi R} (1 + \nu) \sqrt{\frac{1 - \lambda}{\lambda\nu(1 + \lambda)}} , \quad (3.5)$$

$$\Omega_h = \frac{1}{R} \sqrt{\frac{\lambda - \nu}{\lambda(1 + \lambda)}} . \quad (3.6)$$

Here R describes the scale for the solution, and λ and ν are parameters that determine the shape and rotation velocity of the ring[34]. The validity of first law $TdS = dM - \Omega_h dJ$ also requires that λ and ν satisfy the relation (3.1). As a result, a neutral black ring parametrized by M and J can be equivalently described by R and ν . Furthermore, according to [37], a scalar field φ in the neutral black ring reduces to a (1+1)-dimensional free field in the near horizon limit. So the tunneling formalism beyond semiclassical approximation can also be applied to neutral black ring. Now let us turn to the (1+1)-dimensional metric

$$ds^2 = -f(y)dt^2 + \frac{1}{f(y)}dy^2 , \quad (3.7)$$

where,

$$f(y) = \frac{\sqrt{F(y)}}{CR(1+y)} G(y) .$$

Following the standard treatment, consider a massless particle in spacetime (3.7) described by the Klein-Gordon equation

$$-\frac{\hbar^2}{\sqrt{-g}} \partial_\mu [g^{\mu\nu} \sqrt{-g} \partial_\nu] \phi = 0 . \quad (3.8)$$

The semiclassical wave function satisfying the above equation is obtained by making the standard WKB ansatz for ϕ

$$\phi(y, t) = \exp \left[-\frac{i}{\hbar} \mathcal{S}(y, t) \right] . \quad (3.9)$$

Substitute it in (3.8), and expanding the action $\mathcal{S}(y, t)$ in the powers of $\hbar$

$$\mathcal{S}(y, t) = \mathcal{S}_0(y, t) + \sum_i \hbar^i \mathcal{S}_i(y, t) .$$

Following the similar treatment of Kerr-Newman black hole in [28], the final result of the corrected Hawking temperature is

$$T_{cr} = T_{BR} \left(1 + \sum \beta_i \left(\frac{\hbar}{H_{BR}} \right)^i \right)^{-1} . \quad (3.10)$$

Derivation of (3.10) via fermion tunneling is also available, similar to the treatment of other black hole models.

In the context, we have stated that the thermodynamics of neutral black ring can be equivalently parametrized by M and J , or by R and ν . The neutral black ring satisfies the Bekenstein-Hawking law, therefore $\mathcal{A} = \frac{A_h}{4}$. Following the analysis in the previous section, we can prove that $H_{br} = H(R, \nu) = H(\mathcal{A})$. Finally, integrating an equation similar as (2.19), the corrected entropy can be obtained

$$S_{cr} = S_{BR} + \alpha \ln S_{BR} + \cdots + const . \quad (3.11)$$

Now based on the tunneling formulism beyond semiclassical approximation and the first law of thermodynamics, we have shown that the leading order entropy correction of neutral black ring due to quantum effects is also a logarithmic term.

4. Summary

The standard entropy of black holes and black ring entropy should undergoes corrections due to quantum effects. The tunneling formulism beyond semiclassical approximation provides a useful method to calculate the corrected Hawking temperatures and entropy in various models. The entropy corrections obtained using this method have a logarithmic leading order term. This is a natural result from the corrected temperature and the first law of thermodynamics, and coincides with some result obtained by counting the number of microstates in string theory and loop quantum gravity. Maybe this is a somewhat universal phenomenon. In fact, for a black hole model, if the quantum tunneling beyond semiclassical approximation can be used and the corrected temperature can be expressed as (1.2), the leading order correction to its entropy should undergo a logarithmic form, according to the analysis in this paper. However, the exact expressions of black hole and black ring entropy can only be determined in a well-defined quantum gravity theory. We hope that our discussion would be helpful for further understanding of black hole entropy.

Acknowledgments

We thank C.Cao, Q.J.Cao, Y.J.Du, and Q.Ma for useful discussions. The work is supported in part by the NNSF of China Grant No. 90503009, No. 10775116, and 973 Program Grant No. 2005CB724508.

References

- [1] J. M. Bardeen, B. Carter, and S. W. Hawking, *The Four laws of black hole mechanics*, *Commun. Math. Phys.* **31** (1973) 161–170.
- [2] J. D. Bekenstein, *Black holes and entropy*, *Phys. Rev.* **D7** (1973) 2333–2346.
- [3] S. W. Hawking, *Particle Creation by Black Holes*, *Commun. Math. Phys.* **43** (1975) 199–220.
- [4] D. N. Page, *Hawking radiation and black hole thermodynamics*, *New J. Phys.* **7** (2005) 203, [[hep-th/0409024](#)].
- [5] D. V. Fursaev, *Temperature and entropy of a quantum black hole and conformal anomaly*, *Phys. Rev.* **D51** (1995) 5352–5355, [[hep-th/9412161](#)].
- [6] R. B. Mann and S. N. Solodukhin, *Universality of quantum entropy for extreme black holes*, *Nucl. Phys.* **B523** (1998) 293–307, [[hep-th/9709064](#)].
- [7] R. K. Kaul and P. Majumdar, *Logarithmic correction to the Bekenstein-Hawking entropy*, *Phys. Rev. Lett.* **84** (2000) 5255–5257, [[gr-qc/0002040](#)].
- [8] S. Kloster, J. Brannlund, and A. DeBenedictis, *Phase-space and Black Hole Entropy of Higher Genus Horizons in Loop Quantum Gravity*, *Class. Quant. Grav.* **25** (2008) 065008, [[gr-qc/0702036](#)].
- [9] S. Das, P. Majumdar, and R. K. Bhaduri, *General logarithmic corrections to black hole entropy*, *Class. Quant. Grav.* **19** (2002) 2355–2368, [[hep-th/0111001](#)].
- [10] S. S. More, *Higher order corrections to black hole entropy*, *Class. Quant. Grav.* **22** (2005) 4129–4140, [[gr-qc/0410071](#)].
- [11] S. Mukherji and S. S. Pal, *Logarithmic corrections to black hole entropy and AdS/CFT correspondence*, *JHEP* **05** (2002) 026, [[hep-th/0205164](#)].
- [12] S. Carlip, *Logarithmic corrections to black hole entropy from the Cardy formula*, *Class. Quant. Grav.* **17** (2000) 4175–4186, [[gr-qc/0005017](#)].
- [13] M. R. Setare, *Logarithmic correction to the Cardy-Verlinde formula in Achucarro-Ortiz black hole*, *Eur. Phys. J.* **C33** (2004) 555–559, [[hep-th/0309134](#)].
- [14] A. Strominger and C. Vafa, *Microscopic Origin of the Bekenstein-Hawking Entropy*, *Phys. Lett.* **B379** (1996) 99–104, [[hep-th/9601029](#)].
- [15] S. N. Solodukhin, *Entropy of Schwarzschild black hole and string-black hole correspondence*, *Phys. Rev.* **D57** (1998) 2410–2414, [[hep-th/9701106](#)].
- [16] C. Rovelli, *Black hole entropy from loop quantum gravity*, *Phys. Rev. Lett.* **77** (1996) 3288–3291, [[gr-qc/9603063](#)].
- [17] K. A. Meissner, *Black hole entropy in loop quantum gravity*, *Class. Quant. Grav.* **21** (2004) 5245–5252, [[gr-qc/0407052](#)].

- [18] J. B. Hartle and S. W. Hawking, *Path Integral Derivation of Black Hole Radiance*, *Phys. Rev.* **D13** (1976) 2188–2203.
- [19] G. W. Gibbons and S. W. Hawking, *Action Integrals and Partition Functions in Quantum Gravity*, *Phys. Rev.* **D15** (1977) 2752–2756.
- [20] S. M. Christensen and S. A. Fulling, *Trace Anomalies and the Hawking Effect*, *Phys. Rev.* **D15** (1977) 2088–2104.
- [21] K. Srinivasan and T. Padmanabhan, *Particle production and complex path analysis*, *Phys. Rev.* **D60** (1999) 024007, [[gr-qc/9812028](#)].
- [22] M. K. Parikh and F. Wilczek, *Hawking radiation as tunneling*, *Phys. Rev. Lett.* **85** (2000) 5042–5045, [[hep-th/9907001](#)].
- [23] M. K. Parikh, *A secret tunnel through the horizon*, *Int. J. Mod. Phys.* **D13** (2004) 2351–2354, [[hep-th/0405160](#)].
- [24] S. Shankaranarayanan, K. Srinivasan, and T. Padmanabhan, *Method of complex paths and general covariance of Hawking radiation*, *Mod. Phys. Lett.* **A16** (2001) 571–578, [[gr-qc/0007022](#)].
- [25] S. Shankaranarayanan, T. Padmanabhan, and K. Srinivasan, *Hawking radiation in different coordinate settings: Complex paths approach*, *Class. Quant. Grav.* **19** (2002) 2671–2688, [[gr-qc/0010042](#)].
- [26] R. Banerjee and B. R. Majhi, *Quantum Tunneling Beyond Semiclassical Approximation*, *JHEP* **06** (2008) 095, [[arXiv:0805.2220](#)].
- [27] B. R. Majhi, *Fermion Tunneling Beyond Semiclassical Approximation*, *Phys. Rev.* **D79** (2009) 044005, [[arXiv:0809.1508](#)].
- [28] R. Banerjee and S. K. Modak, *Exact Differential and Corrected Area Law for Stationary Black Holes in Tunneling Method*, *JHEP* **05** (2009) 063, [[arXiv:0903.3321](#)].
- [29] S. K. Modak, *Corrected entropy of BTZ black hole in tunneling approach*, *Phys. Lett.* **B671** (2009) 167–173, [[arXiv:0807.0959](#)].
- [30] R. Banerjee, B. R. Majhi, and D. Roy, *Corrections to Unruh effect in tunneling formalism and mapping with Hawking effect*, [arXiv:0901.0466](#).
- [31] R. Banerjee and S. K. Modak, *Quantum Tunneling, Blackbody Spectrum and Non-Logarithmic Entropy Correction for Lovelock Black Holes*, *JHEP* **11** (2009) 073, [[arXiv:0908.2346](#)].
- [32] T. Zhu, J.-R. Ren, and M.-F. Li, *Corrected entropy of high dimensional black holes*, [arXiv:0906.4194](#).
- [33] T. Zhu, J.-R. Ren, and M.-F. Li, *Corrected Entropy of Friedmann-Robertson-Walker Universe in Tunneling Method*, *JCAP* **0908** (2009) 010, [[arXiv:0905.1838](#)].
- [34] R. Emparan and H. S. Reall, *Black rings*, *Class. Quant. Grav.* **23** (2006) R169, [[hep-th/0608012](#)].
- [35] L. Zhao, *Tunnelling through black rings*, *Commun. Theor. Phys.* **47** (2007) 835–842, [[hep-th/0602065](#)].
- [36] Q.-Q. Jiang, *Dirac particles’ tunnelling from black rings*, *Phys. Rev.* **D78** (2008) 044009, [[arXiv:0807.1358](#)].

- [37] B. Chen and W. He, *Hawking Radiation of Black Rings from Anomalies*, *Class. Quant. Grav.* **25** (2008) 135011, [[arXiv:0705.2984](#)].
- [38] M. M. Caldarelli, G. Cognola, and D. Klemm, *Thermodynamics of Kerr-Newman-AdS black holes and conformal field theories*, *Class. Quant. Grav.* **17** (2000) 399–420, [[hep-th/9908022](#)].